

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS ✓
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q	A2	(11) International Publication Number: WO 98/58075 (43) International Publication Date: 23 December 1998 (23.12.98)
(21) International Application Number: PCT/US98/12728 (22) International Filing Date: 17 June 1998 (17.06.98) (30) Priority Data: 60/050,093 18 June 1997 (18.06.97) US Not furnished 17 June 1998 (17.06.98) US (71) Applicants: MONTANA STATE UNIVERSITY [US/US]; Bozeman, MT 59717-3980 (US). UNIVERSITY OF IOWA RESEARCH FOUNDATION [US/US]; Oakdale Research Campus, 100 Oakdale Campus #214 TIC, Iowa City, IA 52242-5000 (US). UNIVERSITY OF ROCHESTER [US/US]; Rochester, NY 14642 (US). (72) Inventors: DAVIES, David, G.; Bozeman, MT (US). COSTERTON, John, William; Bozeman, MT (US). PARSEK, Matthew, R.; - (US). GREENBERG, Everett, P.; - (US). PEARSON, James, P.; - (US). IGLEWSKI, Barbara, H.; - (US). (74) Agents: HANLEY, Elizabeth, A. et al.; Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109 (US).		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: HOMOSERINE LACTONES BIOFILM REGULATING COMPOUNDS AND THEIR METHODS OF USE (57) Abstract <p>A method to control the formulation, persistence and dispersion of microbial biofilms by taking advantage of the natural process of cell-cell communication in bacteria. Addition of N-(3-oxododecanoyl)-L-homoserine lactone (OddHL) and N-butyryl-L-homoserine lactone (BHL) either in combination or separately or the addition of chemicals which will enhance or inhibit the activity of OddHL and BHL can inhibit the formation, persistence or dispersion of bacterial and algal biofilms, in industrial, medical and environmental situations.</p> <div data-bbox="803 1165 1372 1942"></div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

- 1 -

HOMOSERINE LACTONES BIOFILM REGULATING COMPOUNDS AND THEIR METHODS OF USE

Technical Field

5 The present invention relates to homoserine lactone compounds and compositions and their methods of use in biofilm regulation in medical, dental, industrial and environmental settings.

Background

10 Modern methods of direct observation of living biofilms (Lawrence et al., 1993) have established the very complicated structural architecture (Costerton et al., 1995) of these sessile microbial populations. The revelation that biofilms contain distinct microcolonies, separated by discrete water channels (DeBeer et al., 1994), suggested the operation of a cell-cell signaling mechanism that would be sufficient, at a minimum to
15 maintain the patency of these water-filled spaces.

 Prior to 1981, microbiologists had generally assumed that bacteria had neither the requirement nor the capability of producing, cell-cell signaling molecules.

 In 1981, it was shown by Eberhard et al. that the bacterium *Photobacterium fischeri* produces a compound 3-oxo-N-(tetrahydro-2-oxo-3-furanyl) hexanamide
20 also known as vibrio (photobacterium) autoinducer (VAI), which is associated with bacterial luminescence under conditions of high cell density. The cell membrane of *P. fischeri* was shown to be permeable to VAI by Kaplan and Greenberg in 1985. At low bacterial cell densities in broth medium, VAI passively diffuses out of the cells along a concentration gradient, where it accumulates in the surrounding medium. At
25 high cell densities the concentration of VAI outside the cells is equivalent to the concentration of VAI inside the cells. Under such conditions VAI was shown to diffuse back into the cells, resulting in the initiation of transcription of luminescence genes. Using such a system, bacteria are able to monitor their own population density and regulate the activity of specific genes at the population level.

30 For several years it was presumed that the autoinducer involved in bacterial luminescence was unique to the few bacteria that produce light in the marine environment. Then, in 1992, the terrestrial bacterium *Erwinia carotovora* was shown to use an autoinducer system to regulate the production of the B-lactam antibiotic carbapenem (Bainton et al. 1992b). The molecule found to be responsible for
35 autoinduction of carbapenem was shown to be an acylated homoserine lactone (HSL), a member of the same class of molecule responsible for autoinduction in bioluminescence. This finding led to a general search for HSLs in a wide range of bacteria. To affect the

- 2 -

search, a bioluminescence sensor system was developed and used to screen for HSL production in the spent supernatant liquids of a number of bacterial cultures. Many different organisms were shown by the screening to produce HSLs. These included: *Pseudomonas aeruginosa*, *Serratia marcescens*, *Erwinia herbicola*, *Citrobacter freundii*,
5 *Enterobacter agglomerans* and *Proteus mirabilis* (Brainton et al., 1992a; Swift et al. 1993). More recently, the list has grown to include *Erwinia stewartii* (Beck 1993), *Yersinia enterocolitica* (Throup et al., 1995), *Agrobacterium tumefaciens* (Zhang et al., 1993), *Chromobacterium violaceum* (Winston et al., 1994), *Rhizobium leguminosarium* (Schripsema et al., 1996) and others. Today it is generally assumed that all enteric
10 bacteria and the gram negative bacteria generally, are capable of cell density regulation using HSL autoinducers.

In 1993 Gambello et al. (1993) showed that the a-HSL product of the LasI gene of *Pseudomonas aeruginosa* controls the production of exotoxin A, and of other virulence factors, in a cell density dependent manner. Since that time, the production of a
15 large number of *Pseudomonas* virulence factors have been shown to be controlled by a-HSL compounds produced by the LasI and Rhl I regulatory systems (Ochener et al., 1994; Winson et al., 1995; Latifi et al., 1995), in a manner reminiscent of the Lux system. Latifi et. al. (1996) have also shown that many stationary phase properties of *P. aeruginosa*, including those controlled by the stationary phase sigma factor (RpoS), are
20 under the hierarchical control of the LasI and Rhl cell-cell signaling systems. Williams and Brown (1992) have suggested that many of the properties of biofilm bacteria, including their remarkable resistance to antibiotics (Nickel et al., 1985), may derive from the fact that some of their component cells exhibit characteristics of stationary phase planktonic cells.

In all cases, homoserine lactone autoinducers are known to bind to a DNA binding protein homologous to LuxR in *Photobacterium fischeri*, causing a conformational change in activation. This process couples the expression of specific genes to bacterial cell density (Latifi et al., 1996). Regulation of this type been called 'quorum sensing' because it suggests the requirement for a 'quorate' population of
30 bacterial cells prior to activation of the target genes (Fuqua et al., 1994). Expression of certain of these 'virulence factors' has been correlated with bacterial cell density (Finley and Falkow, 1989).

In *P. aeruginosa*, quorum sensing has been shown to be involved in the regulation of a large number of exoproducts including elastase, alkaline protease, LasA protease, hemolysin, cyanide, pyocyanin and rhamnolipid (Gambello et al., 1993; Latifi
35 et al., 1995; Winson et al., 1995; Ochener et al. 1994); but has never before been shown

- 3 -

to be involved in biofilm formation. Most of these exoproducts are synthesized and exported maximally as *P. aeruginosa* enters stationary phase.

It is during stationary phase also, that gram negative bacteria have been shown to develop stress response resistance that is coordinately regulated through the induction of a stationary-phase sigma factor known as RpoS (Hengge-Aronis, 1993). Biofilm bacteria are generally considered to show physiological similarity to stationary phase bacteria in batch cultures. Thus, it is presumed that the synthesis and export of stationaryphase autoinducer-mediated exoproducts occurs generally within biofilms. The stationary phase behavior of biofilm bacteria may be explained by the activity of accumulated HSL within cell clusters. The mechanism causing biofilm bacteria to demonstrate stationary-phase behavior is hinted at by the recent discovery that RpoS is produced in response to accumulation of BHL in *P. aeruginosa* cultures (Latifi et al., 1996).

Biofilms are biological films that develop and persist at interfaces in aqueous environments (Geesey et al., 1977; 1994; Boivin et al., 1991; Khoury et al., 1992; Costerton et al. 1994), especially along the inner walls of conduit material in industrial facilities, in household plumbing systems, on medical implants, or as foci of chronic infections. These biological films are composed of microorganisms embedded in an organic gelatinous structure composed of one or more matrix polymers which are secreted by the resident microorganisms. Biofilms can develop into macroscopic structures several millimeters or centimeters in thickness and can cover large surface areas. These biological formations can play a role in restricting or entirely blocking flow in plumbing systems and often decrease the life of materials through corrosive action mediated by the embedded bacteria. Biofilms are also capable of trapping nutrients and particulates that can contribute to their enhanced development and stability.

The involvement of extracellular polymers in bacterial biofilms has been documented for both aquatic (Jones et al., 1969; Sutherland, 1980) and marine bacteria (Floodgate, 1972), and the association of exopolysaccharides with attached bacteria has been demonstrated using electron, microscopy (Geesey et al., 1977; Dempsey, 1981) and light microscopy (Zobell, 1943; Allison and Sutherland, 1984). The presence of such exopolysaccharides is believed to be involved in the development of the microbial biofilm (Wardell et al., 1983; Allison and Sutherland 1987). Analysis of biofilm bacteria isolated from freshwater and marine environments has shown that the polymers they produce are composed largely of acidic polysaccharides (Fletcher, 1980; Sutherland, 1980; Christensen and Charaklis, 1990). The control and removal of biofilm material from pipe and conduit surfaces has historically been carried out by the addition of corrosive chemicals such as chlorine or strong alkali solutions or through mechanical means. Such treatments are generally harsh to both the plumbing systems and the

- 4 -

environment, and have been necessary due to the recalcitrant nature of biofilms within those systems. The resistance to treatment by biocides has been due in large measure to the protective character of intact biofilm matrix polymers (Srinivasan et al., 1995; Stewart, 1994; Tashiro et al., 1991). In medicine, the use of elevated doses of antibiotics
5 has been necessary in treatment when biofilms are believed to be involved. This is due at least in part to the enhanced protection of biofilm bacteria by the exopolysaccharide matrix material outside the cells (Costerton et al., 1987; Nichols et al., 1989; Anwar et al., 1989).

There is a need in the medical, environmental and industrial arts for the control
10 of biofilm formation. The control of biofilms can be carried out more effectively if the production and regulation of exopolysaccharide material produced by the bacteria can be influenced externally. The present invention overcomes the deficiencies of prior art disinfectant and cleaning compositions by providing a method whereby cell-cell communication in bacteria via LuxR/LuxI homologous systems are manipulated to
15 control biofilm architecture and structural integrity.

Summary of the Invention

The present invention provides a method of regulating biofilm development comprising administering a composition comprising a homoserine lactone compound
20 selected from the group consisting of N-(3-oxododecanoyl) L-homoserine lactone or its analogs and blocking compounds and butyryl L-homoserine lactone or its analogs and blocking compounds.

Also provided is a method for prevention of biofilm matrix polymer development comprising administering a homoserine lactone compound selected from the group
25 consisting of butyryl L-homoserine lactone or its analogs.

The invention advantageously provides a method of biofilm dispersion comprising administering a homoserine lactone compound selected from the group consisting of include L-homoserine lactone or their analogs.

Similarly the invention provides a method of prevention of biofilm dispersion
30 comprising administering an effective amount of a composition including N-(3-oxododecanoyl) L-homoserine lactone or its analogs.

An application of the invention provides a topical dressing for burn patients comprising a homoserine lactone compound selected from the group consisting of N-3(oxododecanoyl)-HSL, include L-homoserine lactone or their analogs.

35 In an alternative embodiment the invention provides a dentifrice or a mouthwash comprising a homoserine lactone compound selected from the group consisting of N-3(oxododecanoyl)-HSL, include L-homoserine lactone or its analogs.

- 5 -

Similarly the invention provides a method for treatment and prevention of dental caries comprising administering composition comprising an effective amount of homoserine lactone compound selected from the group consisting of N-3(oxododecanoyl)-HSL, include L-homoserine lactone or their analogs.

5 Alternatively the invention provides a method for the treatment of acne comprising administering a composition comprising an effective amount of homoserine lactone compound selected from the group consisting of N-3(oxododecanoyl)-HSL, include L-homoserine lactone or its analogs.

Another application of the invention is a method for cleaning and disinfecting
10 contact lenses comprising administering a composition comprising an effective amount of homoserine lactone compound selected from the group consisting of N-3(oxododecanoyl)-HSL, include Lhomoserine lactone or their analogs.

Also provided is a method of treating/(disinfecting and cleaning) medical indwelling devices comprising administering a composition comprising an effective
15 amount of homoserine lactone compound selected from the group consisting of butyryl L-homoserine lactone or its analogs or an N-(3-oxodo-decanoyl) L-homoserine lactone blocking compound. These devices may advantageously include, for example, any indwelling device, for example catheters, orthopedic devices and implants.

The invention provides for a microbial disinfectant composition comprising an
20 effective amount of homoserine lactone compound selected from the group consisting of butyryl L-homoserine lactone or its analogs. In a preferred embodiment the microbe to be disinfected is selected from fungi, protozoa and gram negative bacteria.

The invention provides a method of blocking the binding interaction of homoserine lactones with their corresponding DNA binding protein receptors of LasI
25 and LasR comprising administering a homoserine lactone blocking compound, wherein said blocking interaction prevents biofilm formation.

In still another embodiment the invention provides a method of preventing fouling comprising administering a composition comprising a homoserine lactone compound selected from the group consisting of N-(3-oxododecanoyl) L-homoserine
30 lactone blocking compounds and butyryl L-homoserine lactone or its analogs. In a preferred embodiment the fouling is in a system selected from injection wells for oil recovery, cooling towers, water purification systems, porous media (soil, sand), marine environments and hospital or automotive airconditioning systems.

The invention, advantageously provides a method for the dispersal of
35 environmental concentrations of bacteria comprising administering a homoserine lactone compound selected from the group consisting of N-(3-oxododecanoyl) L-homoserine

- 6 -

lactone blocking compounds and butyryl L-homoserine lactone or its analogs to the environmental concentration of bacteria.

In another application the invention provides a disinfectant coating comprising a homoserine lactone compound selected from the group consisting of

5 N-(3-oxododecanoyl) L-homoserine lactone blocking compounds and butyryl L-homoserine lactone or its analogs copolymerized to a polymer. In a preferred embodiment the coating is applied to drains, shower curtains, grout, toilets, flooring.

In still another embodiment the invention provides for a method of enhancing biofilm formation in fermentation comprising adding N-(3-oxododecanoyl)

10 L-homoserine lactone or its analogs to a fermentation system.

In a medical application the invention provides for a specific therapy to reduce gram negative bacterial without depleting beneficial gram positive bacteria comprising N-(3-oxododecanoyl) L-homoserine lactone blocking compounds and butyryl L-homoserine lactone or their analogs. This method is preferably used in the treatment

15 of a condition selected from urethral e. coli infections and vaginal candida infections.

Alternatively the invention provides a method of treating a condition selected from the group consisting of middle ear infections (children), osteomyelitis and prostatitis comprising administering a homoserine lactone compound selected from the group consisting of N-(3-oxododecanoyl) L-homoserine lactone blocking compounds

20 and butyryl L-homoserine lactone or its analogs, wherein said homoserine lactone compound disperses bacteria making them more susceptible to antibiotics.

Finally, the invention provides for a cleaning composition for cleaning surfaces, for example hard surfaces, woven or unwoven surfaces. Examples of surfaces which may be cleaned and/or cleaning compositions of the invention include toilet bowls, bath

25 tubs, drains, high chairs, countertops (such as those exposed to meats such as chicken), vegetables, meat processing rooms, butcher shops, airducts, airconditioners, carpets, paper or woven product treatment, diapers and healthy air machines. The cleaning product may be in the form of a toilet drop-in for prevention and removal of soil and under rim cleaner for toilets. The compound of the invention may be incorporated into

30 personal care products and skin care products. The compounds of the invention may be used in the preparation of epidermal bandages and lotions. In an alternative embodiment, the compounds of the invention may be incorporated into, for example, after shaves or lotions for the treatment of razor bumps.

The above and other objects of the invention will become readily apparent to

35 those of skill in the relevant art from the following detailed description and figures, wherein only the preferred embodiments of the invention are shown and described, simply by way of illustration of the best mode of carrying out the invention. As is

- 7 -

readily recognized the invention is capable of modifications within the skill of the relevant art without departing from the spirit and scope of the invention.

Brief Description of Drawings

5 Figure 1, shows the mechanism of quorum sensing in planktonic bacteria. Figure 1A, shows that bacteria in both culture elaborate OdDHL (dark circle), and BHL (light circle) into the surrounding medium. At low cell concentrations these HSL molecules passively diffuse across the cell envelop and away from the bacteria. Figure 1B, shows that under conditions of high cell density, these HSL molecules have accumulated to a
10 high concentration and are able to remain within or re-enter the cells, where they bind to cognate receptor LuxR homologous proteins. Once these proteins are bound to the appropriate HSL they in turn bind to regulatory sequences on the chromosome turning on specific genes which produce products such as enzymes, toxins and surfactants.

 Figure 2, shows the structural components of a normal biofilm.

15 Figure 3A, shows the effect of blocking the activity of OdDHL and BHL. When these homoserine lactones are blocked in *P. aeruginosa* the bacteria do not form normal biofilms and they lack the normal architecture associated with these biofilms. Such biofilms can be easily dispersed by the addition of 0.2% sodium dodecyl sulfate. Figure 3B, shows the effect of blocking the activity of BHL in a *P. aeruginosa* biofilm. Under
20 such circumstances, the bacteria elaborate OdDHL (light circles) form a normal looking biofilm but are incapable of undergoing natural detachment since the activity of BHL is associated with detachment. Figure 3C, shows that in a nonnal *P. aeruginosa* biofilm, both homoserine lactones are produced (light and dark circles) and elaborated into the interstices of the exopolymer matrix.

25 Figure 4, shows how biofilms might be manipulated through the use of enhancers or blockers for homoserine lactones. Figure 4A, shows a normal biofilm with bacteria releasing small amounts of both homoserine lactone. Figure 4B, shows how normal detachment occurs in *P. aeruginosa* biofilms. Following the accumulation of BHL, specific enzymes are released from the bacteria which are capable of digesting the
30 extracellular matrix which binds the cells in a biofilm. Once the matrix has been degraded, the cells are free to swim away and disperse from one another. Figure 4C, shows that by the addition of elevated levels of BHL, specific bacteria can be induced to undergo a detachment event artificially.

 Figures 5 photomicrograph in which the mutant defective only in BHL synthesis
35 produced cell clusters that were similar to the wild-type organism.

- 8 -

Figure 6 shows when the double mutant was grown as a biofilm in a medium containing filtered material from the parental wild-type PAO, it developed an intermediate form between the wild-type and the untreated double mutant.

Figure 7 shows the double mutant cultured in medium containing in medium
5 containing a concentration of 10 μ M OdDHL in fresh medium. This resulted in recovery of the intermediary phenotype.

Figure 8 shows, when *P. aeruginosa* PAO1 was treated with 0.2% SDS, no dispersion or release of bacteria from the cell clusters was observed (Figure 8A). When *P. aeruginosa* PAO-JP2 was treated with SDS in the same manner, the cell clusters were
10 shown to disperse completely (Figure 8B), showing a similar effect to what has been seen for *P. aeruginosa* strain 8830 following degradation of alginate.

Figure 9 shows an experiment using the double HSL mutant strain *P. aeruginosa* PAO-JP2, which when grown in medium containing 10 μ M OdDHL was shown to fail to disperse following the addition of sodium dodecyl sulfide (SDS).

Figure 10 shows when the OdDHL mutant *P. aeruginosa* PAO-JP1 was grown as
15 a biofilm, large void spaces were detected in the interior of the cell clusters.

Figure 11 shows an investigation of growing biofilms of *P. aeruginosa* PDO100 and adding BHL to the influent medium at a concentration of 20 μ M after 7 days growth. Following the addition of BHL for 24 hours, no observable effect was detected.
20 Medium flow was then shut off for a period of 16 hours, at which time, significant detachment began to occur and continue for a period of three hours.

Description of the Invention


The inventors have discovered that certain cell communication molecules are
25 responsible for the regulation of microbial biofilm formation, persistence and dispersion. At least two known N-acyl-L homoserine lactones have been found by the inventors to be responsible for the regulation of *Pseudomonas aeruginosa* biofilms; these are, N-(3-oxododecanoyl)-L-homoserine lactone (OdDHL) and N-butyryl-L-homoserine lactone (BHL). The former has been demonstrated to regulate the development of
30 *Pseudomonas aeruginosa* biofilms and to be responsible for maintaining the integrity of biofilm structures by controlling the production of biofilm matrix polymers. The latter has been shown to be involved in the dispersion of *Pseudomonas aeruginosa* biofilms, regulating the production and release of molecules responsible for breaking apart biofilm matrix material. Homoserine lactones have been isolated from a wide range of bacteria
35 and we believe they are responsible for biofilm regulation in organisms other than *P. aeruginosa*.

- 9 -

By artificially manipulating the binding of homoserine lactones to their cognate receptor molecules, the inventors are able to control the formation, development, persistence and dispersion of microbial biofilms. For instance, the addition of an analog which blocks the binding of OdDHL to its cognate receptor (LasR) prevents the
 5 production of matrix polymer material as the bacteria continue to multiply. The result is that cell aggregates formed under these conditions can be easily dispersed by the addition of simple surfactants. Additionally, developed biofilms can be treated with the homoserine lactone BHL to induce the release of enzymes which can digest the biofilm matrix material and disperse the bacteria into the bulk medium. Such treatments could be
 10 used as effective means of controlling biofilm ecology in nature, in industry and in medical applications.


Due to the simple nature of the homoserine lactones as a group, analogs are produced which can act not only, on *P. aeruginosa* but bacteria in general, particularly on pseudomonads and gram negative bacteria. Also, the production of blocking
 15 compounds to the HSLs such as OdDHL and BHL should result in a group of chemicals that can sterically block the binding of HSLs to their cognate receptor molecules, i.e. LasR and RhlR, respectively, and, therefore, block the activity of native HSLs. The simplicity of these molecules also indicates that they should not be antigenic when used as therapeutic agents in the prevention and treatment of bacterial infections. Therefore,
 20 HSLs can be used effectively (alone or in combination with other microbicidal treatments) to treat biofilms in the human body.

N-Butyryl-L-Homoserine Lactone




25

N(3-oxododecanoyl)-L-homoserine Lactone




N-Butyryl-L-Homoserine Lactone compounds



5

N(3-oxododecanoyl)-L-homoserine Lactone compounds



10

wherein R1-R21 is selected from CH3, C1-C4 alkyl group, H, OH, NH2, and SH;

wherein R22 and R23 may be selected from S and O.

wherein R24-R28 is a H or a halogen.

The following are additional analog structures for homoserine lactones of the invention.

15 These are analogs that competitively bind to the cognate receptor protein (LasR or RhlR) and block the binding of OdDHL or BHL, either inducing the receptor protein to bind to its receptor sites on the DNA molecule or preventing the receptor Protein from binding to its receptor sites on the DNA molecule.

1) Alteration of the acyl side chain by increasing or decreasing its length.

20 2) Alteration of the structure of the acyl side chain, such as addition of a double bond or a triple bond between carbon atoms within the acyl side chain.

3) Substitution on carbons in acyl side chain, e.g., the addition of a methyl group or other group such as an oxo-group, a hydroxyl group, an amino group, a sulfur atom or some other atom or R-group to any location along the acyl side chain.

25 4) Substitution on the homoserine lactone ring portion of the molecule. For example: addition of sulfur group to produce a thiolactone.

5) Halogenated acyl furanones have been shown to act as blockers to homoserine lactone cognate receptor proteins.

- 11 -

Also, blocking analogs will bind to the HSL and scavenge it in free form from the environment. Such analogs have a binding site to OdDHL or BHL with a similar structure but greater affinity than the cognate binding protein, LasR or RhIR.

The effects of treating biofilms with homoserine lactones has been demonstrated with *Pseudomonas aeruginosa*. The molecules have generally been isolated from a wide range of bacteria known to be found in biofilms. Among these are the enterobacteria. The presence of the chemicals in a wide range of bacteria indicates that HSLs can be used to effectively treat not only *Pseudomonas sp.* biofilms but also mixed biofilms continuing *Pseudomonas sp.* and biofilms composed of bacteria other than *Pseudomonas aeruginosa*. The application of using HSLs to treat biofilms is, therefore, universal in scope.

Cells of the wild type strain (PAO1) of *Pseudomonas aeruginosa* adhere avidly to a glass surface of the flow cells and, following adhesion, these cells initiate biofilm formation. Direct observations of individual cells containing a Beta-galactosidase reporter gene construct downstream of the Alg C gene (Davies et al., 1995) have shown that alginate production is up regulated within 2-5 minutes following adhesion. The production of this exopolysaccharide matrix material attaches the cells of the wild strain firmly to the substratum, and separates the cells within the developing biofilm until they occupy 15-20% of the volume of the slime-enclosed sessile population. Early in the process of biofilm formation, the sessile cells of the wild (PAO1) strain are seen to become organized into discrete microcolonies (Figure 5) which are separated by well defined water channels (Lawrence et al., 1992). This typical biofilm architecture (Costerton et al., 1995) is detectable by phase microscopy, because the sessile cells are widely separated in the exopolysaccharide matrix (Figure 5A), but it is more clearly seen by confocal scanning laser microscopy (CSLM) (Figure 5B). CSLM of developing biofilms of the wild strain shows the formation of discrete microcolonies, and well defined water channels, and we know that this biofilm architecture is typical of biofilms in general because it is seen in CSLM of mixed species biofilms in natural ecosystems (Costerton et al., 1995).

Cells of mutants of the wild type strain (PAO1) of *Pseudomonas aeruginosa* lack the ability to produce the oxydodecanoyl HSL (PAO-JP1), or lack the ability to produce both the oxydodecanoyl and the butyryl HSL (PAO-JP2), adhere to the glass of the flow cell with an avidity equal to that of the wild type strain. However, after they adhere, they fail to produce alginate. This failure to initiate biofilm formation by alginate production is evidenced by the negative values for uronic acid production by sessile cells of these mutants (Table 2), even after they have been in the adherent state for 7 days.

- 12 -

Phase contrast microscopy shows that the adherent cells of the single (JP1) and double (JP2) mutants are immediately juxtaposed to each other, like cordwood in a pile (Figure 4, Figure 3), and there is no evidence of cell-cell separation by alginate production.

5 The complete failure of both the single mutant, which is incapable of making OdDHL, and the double mutant, which is incapable of making either OdDHL or BHL, to form biofilms is thus unequivocally shown by both chemical and microscopy methods. This failure of the HSL negative mutants to form biofilms, following adhesion, is further evidenced by the fact that their adherent cells are readily removed from the glass surface
10 of the flow cell by simple washing with 0.2% SDS. These data are sufficient to show that the formation of biofilms by cells of *Pseudomonas aeruginosa* is dependent on the production of the acyl homoserine lactone OdDHL. This a-HSL signalling system is hierarchically superior to the sigma factor regulated by AlgT (Deretic et al., 1996), in that alginate is not formed by adherent cells that express AlgT if OdDHL production is
15 deficient.

This control of biofilm formation by OdDHL opens up countless applications for the practical control of biofilm problems in industry, and in medicine (Khoury et al., 1994). The inventors show that analogues of the LasI product (OdDHL) will bind to the LasR DNA binding protein, and block its ability to express the genes that regulate
20 alginate synthesis and biofilm formation.

The a-HSLs are relatively simple molecules (Pearson et al. 1994, Fuqua et al. 1994, Pearson et al. 1995), and several analogues of their basic structures have been produced in several laboratories (Eberhard, et al., 1986). The halogenated furanone that appears to control biofilm formation on the red alga, *Delisea pulchra*, in the marine
25 environment (Givskov et al., 1996), has a molecular structure that is similar to that of the a-HSLs.

In industrial systems in which biofilm formation is beneficial such as commercial bioreactors and fermentation systems, the application of natural a-HSL molecules are effective in enhancing biofilm formation. The inventors have entered a new era of the
30 control of bacterial activities, with the advent of commercial a-HSLs and a-HSL analogues, that parallel modern biocontrol measures in other fields.

The simple maintenance of well defined water channels, throughout the biofilms, virtually requires some form of cell-cell signaling. When planktonic cells of the gram negative bacteria that predominate in biofilms in aquatic systems, notably *Pseudomonas*
35 *aeruginosa*, have been examined in batch culture, several cell-cell signaling mechanisms have been discovered. These signaling mechanisms, some of which have now been described in exquisite molecular detail, control several important aspects of cellular

- 13 -

behavior in these planktonic cells. It was reasoned that the signaling mechanisms that operate in planktonic cells in batch cultures, a form of growth that is very rare in nature, probably actually evolved to control cellular behavior in biofilms in real ecosystems. For this reason the inventors examined the ability of mutants that are unable to synthesize specific signaling molecules, acyl homoserine lactones (a-HSLs), to form biofilms. A mutant (JP1) of the PAO strain of *Pseudomonas aeruginosa*, that is unable to synthesize the oxydodecanoyl HSL, was able to adhere to surfaces in a flowing system, but it was unable to form a biofilm. Adherent cells of this mutant grew on the surface, and formed amorphous masses of cells. They were unable to synthesize the exopolysaccharide (alginate) that forms the matrix of their biofilms, and therefore they were unable to produce the complex of microcolonies and water channels that characterize biofilms of these organisms. Another mutant (PD0100) of the PAO1 strain of *Pseudomonas aeruginosa*, that lacks the ability to synthesize butyryl HSL, was unable to produce the programmed detachment of planktonic cells from mature biofilms. This programmed detachment is a property of the wild type PAO1 strain that is normally triggered by the cessation of flow in a continuous system. The significant role of a-HSL signaling molecules in two important cellular behaviors, *a propos* of biofilm formation by cells of a common biofilm organism, indicates that we have identified a class of cell-cell communication molecules that may be used to control biofilms in many ecosystems. The disinfectant of the invention, in a preferred embodiment is useful as a disinfectant for gram negative bacteria selected from *Pseudomonadaceae*, *Azotobacteraceae*, *Rhizobiaceae*, *Methylococcaceae*, *Halobacteriaceae*, *Acetobacteraceae*, *Legionellaceae*, *Neisseriaceae*, Other Genera.

The following is a list of groups of Gram-Negative bacterial that have members which use homoserine lactones for cell-cell communication: anaerobic Gram Negative Straight, Curved and Helical Rods; *Bacteroidaceae*; The Rickettsias and Chlamydias; Dissimilatory Sulfate - or Sulfur-Reducing Bacteria; the Mycoplasmas; The mycobacteria; Budding and/or Appendaged Bacteria; Sheathed Bacteria; Nocardioforms; and Actinomycetes, for example. See Bergey's Manual of Systematic Bacteriology, First Ed., John G. Holt, Editor in Chief (1984) incorporated herein by reference.

All publications, patent applications, patents, and other references mentioned herein are incorporated by reference. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

Example 1

The most unequivocal experimental design, to determine the role of a-HSL signal molecules on the formation of biofilms by cells of *P. aeruginosa*, was to use direct microscopic methods to monitor biofilm formation by cells of a-HSL negative mutants.

5 For this reason planktonic cells of a wild type strain (PAO1), and of three mutants incapable of synthesizing specific a-HSLs, were introduced into flow cells, and adhesion and biofilm formation were monitored by means of confocal scanning laser microscopy (CSLM). Using these techniques, it is possible to monitor the development of live biofilms of the strains of interest.

10 **Bacteria and media.** The Bacteria strains used in this study are listed in Table 1. All experiments were performed using a defined culture medium containing the following, in grams per liter: sodium lactate, 0.05; sodium succinate, 0.05; ammonium nitrate, 0.05; KH₂PO₄, 0.19; K₂HPO₄ 0.63; Hutner Salts (Cohen-Bazire, 1957), 0.01; glucose, 1.0; and L-histidine, 0.01. Solid R2A medium was used for the enumeration of
15 bacteria from continuous culture experiments. HgCl₂ (7.5 ug/ml in continuous culture and 15 ug/ml on solid medium) and tetracycline 25 ug/ml in continuous culture and 50 ug/ml on solid medium) were used to ensure plasmid and transposon maintenance during experiments.

Table 1

Bacterial strains and plasmids used in this work

<i>P. aeruginosa</i>	Relevant Characteristics	Source/Reference
PAO1	wild type	Holloway (1995)
PAO = JP1	<i>las::tet</i> strain PAO1 derivative	Pearson et al. (1977)
PDO100	<i>rhlI::Tn501-2</i> strain PAO1 derivative	Brint and Ohman (1995)
PAO-JP2	<i>lasI::, rhlITb501-2</i> PAO1 derivative	Pearson et al. (1997)

Continuous culture studies. A continuous culture apparatus was developed to observe the growth and development of biofilms attached to a glass substratum (Figure 8). The apparatus was configured as a once through flow cell system. The influent defined culture medium was retained in a four liter glass reservoir. Medium from the influent reservoir was pumped through silicone tubing via a Masterflex pump to an aeration flask sparked with a filtered air. The aerated medium was pumped to a flat plate flow cell using a Masterflex 8 roller-head peristaltic pump at a flow rate of 0.13 ml min⁻¹. The flow cell was constructed of polycarbonate having a depth of 1.0 mm, a width of 1.4 cm, and a length of 4.0 cm, the upper face was capped with a glass coverslip. The
25
30

- 15 -

glass coverslip was used as a substratum for bacterial attachment and biofilm development because it is relatively inert material and is transparent. Flow through the cell was laminar, having a Reynolds number of 0.17, with a fluid residence time of 0.43 min. The flow cell was sealed to prevent contamination and affixed to the stage of an Olympus BH2 microscope. Medium leaving the flow cell was pumped to an effluent reservoir via silicone tubing. The entire system was closed to the outside environment but maintained in equilibrium with atmospheric pressure by a 0.2 μ m pore size gas permeable filter fitted to each flask.

Log phase *P. aeruginosa* were inoculated through a septum approximately 1 cm upstream from the flow cell while flow was maintained. Bacteria were allowed to attach to and grow on the surfaces of the system downstream from the site of inoculation over a period of 24 h. Flow through the system was then increased to remove any bacteria attached to the inside surface of the coverslip (as determined by microscopy). Bacteria shed from biofilm upstream from the flow cell were then allowed to recolonize the surfaces of the flow cell under conditions of normal flow. Cells attached to the inner surface of the glass coverslip were viewed by transmitted light using a 40 x magnification A40PL and a 50x magnification ULWD MSPlan long working distance Olympus objective lens to detect total cells. All images were stored as separate files for subsequent retrieval and analysis.

Biofilm development. measurement of cell cluster size, thickness. Description of cluster morphology (core, channel, streamer). Modified Lowry protein assay. The Lowry protein assay was performed on samples as described previously (Peterson, 1977) and analyzed with a Milton Roy Spectronic 601 spectrophotometer.

Uronic acid assay. Total uronic acids were measured in thawed samples of scraped biofilm and whole culture following the method of Kintner and Van Buren (1982) using a Milton Roy Spectronic 601 spectrophotometer. A total polysaccharide assay and lipopolysaccharide analysis were also performed.

Biofilm architecture. The growth and development of biofilms has been shown to result in the production of specific architectural components (Costerton et al. 1995). The inventors show that biofilm architecture is influenced by cell-cell communication.

Biofilms of *P. aeruginosa* PAO1 were grown in a bioreactor and examined by microscopy coupled with image analysis. Development of the wild-type organism into a mature biofilm over a two week period, resulted in cell clusters ranging in size from 40-120 μ m with an average thickness of (102.3 μ m sd=20.5 n=20). These cell clusters were shown to contain water channels, have few cells attached to the substratum and were composed of bacteria well separated from one another (Figure 5 - confocal image composite). These cell clusters were then compared with those developed by *P.*

- 16 -

aeruginosa which were defective in the ability to synthesize the homoserine lactone molecules OdDHL, BHL or both. Under identical experimental conditions, cell clusters developed by the mutant strain *P. aeruginosa* JP2 which lacked the ability to synthesize either of the homoserine lactones, the architectural components of the wild-type organism were found to be missing (Figure 6). The clusters ranged in size from 20-40 um with an average thickness of 23.5 um sd= 9.8 n=20. Cells in these clusters were densely packed and did not develop water channels. When *P. aeruginosa* PAO-JP1, defective only in OdDHL were grown under similar conditions, they were shown to produce clusters similar in size to the *P. aeruginosa* PAO-JP2 mutant (average thickness = 22.8 um, sd=10.0, n=20), but containing large spaces devoid of cells in the cluster interior (Figure 7). The mutant defective only in BHL synthesis produced cell clusters that were similar to the wild-type organism (average thickness =100.1 um, sd=25.2, n=20) (Figures 9).

Example 2

To confirm that homoserine lactone was responsible for the architectural differences noted between wild-type and mutant biofilms, an experiment was performed to demonstrate that addition of filterable material collected from medium in which the wild-type organism had grown would recover the wild-type architecture in the double mutant, *P. aeruginosa* PAO-JP2. When the double mutant was thus grown as a biofilm, it developed an intermediate form between the wild-type and the untreated double mutant (Figure 10). The interior of the cell clusters appeared similar to the untreated *P. aeruginosa* PAO-JP2 and the exterior of the cell clusters appeared similar to the wild-type organism. This experiment was repeated, culturing the double mutant using a concentration of 10 μ M OdDHL in fresh medium. This resulted in recovery of the intermediary phenotype as was observed when the cells were grown in the presence of spent medium (Figure 11). These results indicated that biofilm architecture *P. aeruginosa* PAO1 biofilms is conferred by cell-cell communication. The inventors conclude that OdDHL is able to control this architectural development.

Biofilm matrix polymer. The architectural differences noted when comparing biofilms developed by wild-type and HSL mutant *P. aeruginosa* led us to predict that matrix polymer production and regulation are controlled by homoserine lactone. Biofilm samples of *P. aeruginosa* PAO1 and *P. aeruginosa* PAO-P2 were cultured for two weeks in a biofilm reactor. When these cultures were analyzed for uronic acids production, the wild-type strain was shown to produce detectable levels, however, none were detectable in the double mutant (Table 2).

- 17 -

This result indicated that the strain *P. aeruginosa* PAO-JP2 does not produce detectable alginate in continuous culture. When this strain is cultured in spent medium from the wild-type, filtered and amended with glucose, the production of uronic acids was recovered. No uronic acids were detectable in the filtered medium from the wild-type organism. The uronic acids assay detects mannuronic acid which is found in alginate and certain forms of lipopolysaccharide (LPS). The results indicated that one or both of these compounds is under the regulation of OdDHL.

Table 2.

Uronic Acids Production in *P. aeruginosa* PAO1 biofilms.

Sample	Uronic Acids / Protein (ug/ug)
<i>P. aeruginosa</i> PAO1	$3.97 \times 10^{-4} \pm 0.41 \times 10^{-4}$
<i>P. aeruginosa</i> PAO-JP2	ND ^b
Filtered medium	$1.62 \times 10^{-5} \pm 0.20 \times 10^{-5}$
<i>P. aeruginosa</i> PAO-JP2 ^a	ND ^b

^a Cells grown in filtered medium from *P. aeruginosa* PAO1 culture.

^b Not detectable.

It has been shown that in mucoid strains of *P. aeruginosa*, alginate lyase is capable of degrading extracellular alginate (Boyd et al. 1994). Further studies by the inventors demonstrated that alginate lyase can degrade extracellular alginate when released artificially from *P. aeruginosa* strain 8830 in biofilms. Following the destruction of extracellular alginate, these bacteria can be completely dispersed through the addition of 0.2% sodium dodecyl sulfate (SDS) (Davies, 1996).

In the present study, treatment with the detergent was shown not to affect the biofilm in the absence of released alginate lyase. When *P. aeruginosa* PAO1 was treated with 0.2% SDS under similar experimental conditions, no dispersion or release of bacteria from the cell clusters was observed (Figure 12, panel A). When *P. aeruginosa* PAO-JP2 was treated with SDS in the same manner, the cell clusters were shown to disperse completely (Figure 12, panel B), showing a similar effect to what had been seen for *P. aeruginosa* strain 8830 following degradation of alginate. This experiment was repeated using the single HSL mutant strain *P. aeruginosa* PAO-JP1, which was shown to disperse completely following the addition of SDS (Figure 13). When *P. aeruginosa* PAO-JP1 was grown in the presence of 10 μ M OdDHL, treatment with 0.2% SDS did

- 18 -

not disperse bacteria in the cell clusters. The presence of the homoserine lactone, therefore, was shown to be responsible for resistance to dispersion by detergent action.

Biofilm dispersion. The role of OdDHL has been shown to include regulation of the development of biofilm architecture and resistance to dispersion by detergent. The inventors believe that BHL is involved with natural dispersion of bacteria in biofilms. When the OdDHL mutant *P. aeruginosa* PAO-JP1 was grown as a biofilm, large void spaces were detected in the interior of the cell clusters (Figure 14). During the growth phase of cell clusters of *P. aeruginosa* PAO-JP1, central void spaces developed after 7 days in clusters greater than 50 μm in diameter. These voids had previously been occupied by bacteria that were observed to become actively motile and eventually swim away from the cluster interior via a break through the cluster wall. It was postulated that the presence of such central voids, which were not detected in cell clusters formed by *P. aeruginosa* PAO-JP2 or *P. aeruginosa* PDO100, indicated the possibility that BHL is responsible for the release of enzymes which can degrade matrix polymer material. The inventors investigated this by growing biofilms of *P. aeruginosa* PDO100 and adding BHL to the influent medium at a concentration of 20 μM after 7 days growth. Following the addition of BHL for 24 hours, no observable effect was detected. Medium flow was then shut off for a period of 16 hours, at which time, significant detachment began to occur and continue for a period of three hours (Figure 15). When *P. aeruginosa* PDO100 was grown in the absence of BHL, medium flow was turned off after 7 days. Following cessation of flow, no dispersion of cell clusters was observed over the duration of the observation period of 96 hours.

Example 3

P. aeruginosa PAO1 mutants which are not able to produce the homoserine lactones OdDHL and BHL produce biofilms which lack the complex architecture of wild-type *P. aeruginosa* PAO1 biofilm. The complex architecture of wild-type *P. aeruginosa* biofilms includes: extensive matrix polymer, void spaces, clusters, streamers and minimal spacing between individual organisms both at the substratum and within cell clusters.

P. aeruginosa PAO1 mutants which are not able to produce the homoserine lactones OdDHL and BHL produce biofilms which are completely dispersed into free floating individual cells following treatment with 0.21 SDS. Wild-type *P. aeruginosa* biofilms are not effected by the addition of 0.2% SDS. *P. aeruginosa* PAO1 mutants which are not able to produce the homoserine lactone OdDHL and BHL do not produce detectable amounts of uronic acids. Wild-type *P. aeruginosa* PAO1 produces detectable

- 19 -

uronic acids which are believed to signify the production of alginate (a polyuronic acid) which is generally considered a principal biofilm matrix polymer.

P. aeruginosa PAO1 mutants which are not able to produce the homoserine lactone OdDHL have been shown to produce cell clusters which have huge void spaces in the interior. This phenomenon has been observed in wild-type *P. aeruginosa* PAO1 but not in *P. aeruginosa* PAO1 mutants which do not produce the homoserine lactones OdDHL and DHL.

P. aeruginosa PAO1 mutants which are not able to produce the homoserine lactone BHL do not undergo biofilm dispersion when treated in a manner similar to wild type biofilms. Also, the addition of BHL to these mutant biofilms causes the bacteria to disperse when left under quiescent conditions for up to 18 hours.

Example 4

The compounds of the present invention are useful in pharmaceutical compositions for systemic administration to humans and animals in unit dosage forms, such as tablets, capsules, pills, powders, granules, suppositories, sterile parenteral solutions or suspensions, sterile non-parenteral solutions or suspensions oral solutions or suspensions, oil in water or water in oil emulsions and the like, containing suitable quantities of an active ingredient. Topical application can be in the form of ointments, creams, lotions, jellies, sprays, douches, and the like. For oral administration either solid or fluid unit dosage forms can be prepared with the compounds of the invention. The compounds are useful in pharmaceutical compositions (wt %) of the active ingredient with a carrier or vehicle in the composition in about 1 to 99% and preferably about 5 to 15%.

Either fluid or solid unit dosage forms can be readily prepared for oral administration. For example, the compounds of Formula I can be mixed with conventional ingredients such as dicalciumphosphate, magnesium aluminum silicate, magnesium stearate, calcium sulfate, starch, talc, lactose, acacia, methyl cellulose and functionally similar materials as pharmaceutical excipients or carriers. A sustained release formulation may optionally be used. Capsules may be formulated by mixing the compound with a pharmaceutical diluent which is inert and inserting this mixture into a hard gelatin capsule having the appropriate size. If soft capsules are desired a slurry of the compound with an acceptable vegetable, light petroleum, or other inert oil can be encapsulated by machine into a gelatin capsule.

Suspensions, syrups and elixers may be used for oral administration of fluid unit dosage forms. A fluid preparation including oil may be used for oil soluble forms. A vegetable oil such as corn oil, peanut oil or safflower oil, for example, together with

- 20 -

flavoring agents, sweeteners and any preservatives produces an acceptable fluid preparation. A surfactant may be added to water to form an emulsion for fluid unit dosages. Hydro-alcoholic pharmaceutical preparations may be used having an acceptable sweetener such as sugar, saccharine or a biological sweetener and a flavoring agent in the form of an elixer.

Pharmaceutical compositions for parenteral and suppository administration can also be obtained using techniques standard in the art.

A preferred use of the compounds according to the invention is as topical agents. Another preferred use of the compounds is in a transdermal parenteral anti-inflammatory pharmaceutical preparation, which is particularly useful in the treatment of burns and open wounds. Accordingly, compositions suitable for administration to these areas are particularly included within the invention. The above parenteral solutions or suspensions may be administered transdermally and, if desired a more concentrated slow release form may be administered. Accordingly, incorporation of the active compounds in a slow release matrix may be implemented for administering transdermally. The compounds may be administered transdermally at about 1 to 99% of the composition and preferably about 5 to 15% wt% of the active ingredient in the vehicle or carrier.

Transdermal therapeutic systems are self-contained dosage forms that, when applied to intact skin, deliver drug(s) at a controlled rate to the systemic circulation. Advantages of using the transdermal routing include: enhanced therapeutic efficacy, reduction in the frequency of dosing, reduction of side effects due to optimization of the blood-concentration versus time profile, increased patient compliance due to elimination of multiple dosing schedules, bypassing the hepatic "first-pass" metabolism, avoiding gastrointestinal incompatibilities and providing a predictable and extended duration of activity. However, the main function of the skin is to act as a barrier to entering compounds. As a consequence, transdermal therapy has so far been restricted to a limited number of drugs that possess the desirable physiochemical properties for diffusion across the skin barrier. One effective method of overcoming the barrier function of the skin is to include a penetration enhancer in the formulation of a transdermal therapeutic system. See Barry, Brian W.: Dermatological Formulations: Percutaneous Absorption (Dekker, New York, 1983); Bronough et al, Percutaneous Absorption. Mechanisms-Methodology-Drug Delivery, (Marcel Dekker, New York, NY 1985); and Monkhouse et al, Transdermal drug deliver problems and promises. Drug Dev. Ind. Pharm., 14, 183-209 (1988).

A penetration enhancer is a chemical compound that, when included in a formulation, temporarily increases the permeability of the skin to a drug allowing more of the drug to be absorbed in a shorter period of time. Several different types of

penetration enhancers have been reported such as dimethylsulfoxide, n-decyl methyl sulfoxide, N,N-dimethylacetamide, N,N-dimethylformamide, 1-dodecylazacycloheptan-2-one (Azone), propylene glycol, ethanol, pyrrolidones such as N-methyl-2-pyrrolidone (NMP) and surfactants. See Bronough et al, supra, and
5 Stoughton et al, Azone: a New Non-toxic enhancer of percutaneous penetration. Drug Dev. Inc. Pharm., 9, 725-744 (1983).

N-methyl-2-pyrrolidone is a versatile solvent which is miscible with water, ethyl alcohol, ether, chloroform, benzene, ethyl acetate and carbon disulfide. N-methyl-pyrrolidone has been widely used as a solvent in industrial processes such as petroleum
10 refining, GAF Corp.: "M-Pyrol (N-methyl-2-pyrrolidone) Handbook.", GAF Corp., New York, 1972. It is currently used as a solubilizing agent in topical and parenteral veterinary pharmaceuticals Wells, D.A. et al: Disposition and Metabolism of Double-Labeled [³H and ¹⁴C] N-methyl-2pyrrolidone in the Rat. Drug Met. Dispos., 16, 243-249 (1988). N-methylpyrrolidone has also been shown to be an effective
15 penetration enhancer. Barry et al, Optimization and Bioavailability of Topical Steroids: Penetration Enhancers Under Occlusion. J. Inv. Derm., 82, 49-52 (1984); Akter et al, Absorption Through human Skin of Ibuprofen and Flurbiprofen; Effect of Dose variation, Deposited Drug Films, Occlusion and the Penetration Enhancer N-methyl-2-pyrrolidone. J. Pharm. Pharmacol., 37, 27-37 (1984); Hølegaard et al,
20 Vesical Effect on Topical Drug Delivery IV. Effect of N-methylpyrrolidone and Polar Lipids on Percutaneous Transport. Int. J. Pharm., 43, 233-240 (1988); Sugibayashi et al, Effect of Several Penetration Enhancers on the Percutaneous Absorption of Indomethacin in Hairless Rat. Chem. Pharm. Bull., 36, 1519-1529 (1988); Bennett et al, Optimization of Bioavailability of Topical Steroids: Non-occluded penetration
25 Enhancers Under Thermodynamic Control. J. Pharm. Pharmacol., 37, 298-304 (1985); Sasaki et al, Enhancing Effect of Pyrrolidone Derivatives on Transdermal Drug Delivery. 1. Ing. J. Pharm., 44, 14-24 (1988); Lee et al, Toxicity of N-methyl-2-pyrrolidone (NMP): Tetratogenic, Subchronic and Two-year Inhalation Studies, Fund. Appl. Tox., 9, 222-235 (1987).

30 The above and other drugs can be present in the reservoir alone or in combination form with pharmaceutical carriers. The pharmaceutical carriers acceptable for the purpose of this invention are the art known carriers that do not adversely affect the drug, the host, or the material comprising the drug delivery device. Suitable pharmaceutical carriers include sterile water; saline, dextrose; dextrose in water or
35 saline; condensation products of castor oil and ethylene oxide combining about 30 to about 35 moles of ethylene oxide per mole of castor oil; liquid acid; lower alkanols; oils such as corn oil; peanut oil, sesame oil and the like, with emulsifiers such as mono- or

- 22 -

di-glyceride of a fatty acid, or a phosphatide, e.g., lecithin, and the like; glycols; polyalkylene glycols; aqueous media in the presence of a suspending agent, for example, sodium carboxymethylcellulose; sodium alginate; poly(vinylpyrrolidone); and the like, alone, or with suitable dispensing agents such as lecithin; polyoxyethylene stearate; and
5 the like. The carrier may also contain adjuvants such as preserving stabilizing, wetting, preserving stabilizing, emulsifying agents and the like together with the penetration enhancer of this invention.

The effective dosage for mammals may vary due to such factors as age, weight, activity level or condition of the subject being treated. Typically, an effective dosage of
10 a compound according to the present invention is about 2.5 μM to 50 μM for the OdDHL compound and 5-100 μM for BHL, preferably 10-15 μM , when administered by either oral or rectal dose from 1 to 3 times daily. The required dose is less when administered parenterally, intramuscularly or transdermally, 1 or 2 times a day for an adult human.

15 Compounds of the present invention may be administered topically at about 1 to 99 wt% of the composition, and preferably about 5 to 35 wt%.

Example 5

Disinfectant compositions include the compositions of the invention and
20 preferably include surfactant compositions known in the art. These include but are not limited to anionic surfactants, cationic surfactants, non-ionic surfactants and amphoteric or ampholytic surfactants, including Sodium Dodecyl Sulfate; Quaternary Ammonium Compounds; alkyl pyridinium iodides; Tween 80, Tween 85, Triton X-100; Brij 56; Biological surfactants; Rhamnolipid, Surfactin, and Visconsin, for example.
25 The disinfectant composition may be applied in known areas and surfaces where disinfection is required, including but not limited to drains, shower curtains, grout, toilets and flooring. A particular application is on hospital surfaces and medical instruments.

Example 6

30 The compositions of the invention may be formulated into a copolymer coating to provide disinfectant surfaces on products.

Example 7

A dentifrice or mouthwash containing the compounds of the invention may be formulated by adding the compounds of the invention to known dentifrice and mouthwash formulations as set forth in Remington's Pharmaceutical Sciences, 18th Ed., Mack Publishing Co., 1990, Chapter 109 (incorporated herein by reference in its entirety). The dentifrice may be formulated as a gel, paste, powder or slurry. The dentifrice may include binders, abrasives, flavoring agents, foaming agents and humectants. Mouthwash formulations are known in the art, and the compounds of the invention may be advantageously added to them.

From the above disclosure certain practical applications are clear. The production, dispersion, binding or concentration of OdDHL and BHL in *P. aeruginosa* can be manipulated to interfere with biofilm architecture. Biofilm architecture is believed to be responsible for fluid and chemical transport in biofilms, for resistance to fluid shear and for the integration of multiple species into biofilms. It, therefore, becomes obvious that manipulation of the effects of OdDHL and BHL in *P. aeruginosa* developed and developing biofilms could effect fluid and chemical transport, resistance to fluid shear and the integration of multiple species into biofilms.

By blocking the activity of the homoserine lactones OdDHL and DHL during biofilm development, *P. aeruginosa* biofilms can be dispersed by the addition of 0.2% or higher concentration of SDS. Detergents other than SDS show potential for the same effect. Such a treatment in combination with any microbicidal agent will significantly increase the efficacy of treating biofilms in industry, the environment and in medical applications.

The production of matrix polymer in biofilms not able to produce the homoserine lactones OdDHL and BHL is significantly impaired more recent results indicate that only OdDHL is needed for matrix polymer production. By blocking the production of matrix polymer in biofilms produced by *P. aeruginosa*, the effects of the matrix polymer in these biofilms should be abrogated. These effects include but are not limited to interference with the transport of chemicals and nutrients into and away from cells within the biofilm production against the effects of reactive chemicals, protection from grazing by eukaryotic organisms, ability of biofilm to bind ions (particularly multivalent cations), protection from attack by cells and chemicals of the immune system, protection from variations in pH and possibly electrical and/or magnetic interference with normal biofilm development and persistence.

The addition of BHL should induce a dispersion response in developed biofilms of *P. aeruginosa*. BHL when added to biofilm cells, should induce those cells to produce and/or release enzymes which will digest the biofilm matrix polymer materials

- 24 -

and cause the bacteria within the biofilm to separate from one another. Following dispersion, these bacteria should be easily treated using conventional methods, such as the use of biocides, antibiotics, detergents, radiation, etc. Used by itself or in concert with microbicidal treatments, the application of BHL and/or BHL analogs should be

5 highly effective in destroying developed biofilms or in the prevention of biofilm development. Such treatments should be effective in controlling biofilms in the household, in industry, in the environment and in medicine.

References

- Bainton, N.J., B.W. Bycroft, S.R. Chhabra, P. Stead, L. Gledhill, P.J. Hill, C.E.D. Rees, M.K. Winson, G.P.C. Salmond, G.S.A.B. Stewart and P. Williams. 1992a. A general
5 role for the *lux* autoinducer in bacterial cell signalling: control of antibiotic synthesis in *Erwinia*. *Gene*. 116:87-91.
- Bainton, N.J., P. Stead, S.R. Chhabra, B.W. Bycroft, G.P.C. Salmond, S.A.A.B. Stewart and P. Williams. 1992b. N-(3-oxohexanoyl)-1-homoserine lactone regulates carbapenem
10 antibiotic production in *Erwinia carotovora*. *Biochem. J.* 288:997-1004.
- Beck von B.S. and S.K. Farrand. 1995. Capsular polysaccharide biosynthesis and pathogenicity in *Erwinia carotovora*. *J. Bacteriol.* 177:5000-5008.
- 15 Boyd, A. and A.M. Chakrabarty. 1994. Role of the alginate lyase in cell detachment of *Pseudomonas aeruginosa*. *Appl. Environ. Microbiol.* 60:2325-2359.
- Brint, J.M. and D.E. Qhman. 1995. Synthesis of multiple exoproducts in *Pseudomonas aeruginosa* is under the control of RH1R-Rh1I. Another set of regulators in strain PAO1
20 with homology to the autoinducer-responsive LuxR-LuxI family. *J. Bacteriol.* 177:7155-7163.
- Cohen-Bazire, G., W.R. Sistrom and R.Y. Stanier. 1957. Kinetic studies of pigment synthesis by non-sulfur purple bacteria. *J. Cell. Comp. Physiol.* 49:25-68.
- 25 Davies, D.G. 1996. Regulation of alginate biosynthesis in *Pseudomonas aeruginosa* biofilms. Doctoral Thesis Montana State University, Bozeman, MT.
- Eberhard, A., A.L. Burlingame, C. Eberhard, G.L. Kenyon, K.H. Nealson and N.J. Oppenheimer. 1981. Structural identification of autoinducer of *Photobacterium fischeri*
30 *luciferase*. *Biochemistry* 20:2444-2449.
- Eberhard, A., C.A. Widrig, P. McBath and J.B. Schineller. 1986. Analogs of the autoinducer of bioluminescence in *Vibrio fischeri*. *Arch. Microbiol.* 146:35-40.
- 35 Finlay, B.B. and Falkow, S. 1989. Common themes in microbial pathogenicity. *Microbiol. Rev.* 53:210-230.

- Fuqua, W.C., S.C. Winans and E.P. Greenberg. 1994a. LasR of *Pseudomonas aeruginosa* is a transcriptional activator of the alkaline protease gene (*apr*) and an enhancer of exotoxin A expression. *Infect. Immun.* 61:1180-1184.
- 5 Fuqua, W.C., S.C. Winans and E.P. Greenberg. 1994b. Quorum sensing in bacteria: the LuxR-LuxI family of cell density-responsive transcriptional regulators. *J. Bacteriol.* 176:269-275.
- 10 Givskov, M., R. DeNys, M. Manefield, L. Gram, R. Maximilten, L. Everl, S. Molin, P.D. Steinberg and S. Kjelleberg. 1996. Eukaryotic interference with homoserine lactone-mediated prokaryotic signalling. *J. Bacteriol.* 178:6618-6622.
- 15 Latifi, A., M. Foglino, K. Tanaka, P. Williams and A. Lazdunski. 1996. A hierarchical quorum sensing cascade in *Pseudomonas aeruginosa* links the transcriptional activators. LasR and RhIR (VsmR) to expression of the stationary-phase sigma factor RpoS. *Mol. Microbiol.* 21:1137-1146.
- 20 Kaplan, H.B. and E.P. Greenberg. 1985. Diffusion of autoinducer is involved in regulation of the *Vibrio fischeri* luminescence system. *J. Bacteriol.* 163:1210-1214.
- Kintner, P.K., III and J.P. Van Buren. 1982. Carbohydrate interference and its correction in pectin analysis using the m-hydroxydiphenyl method. *J. Food Sci.* 47:756-760.
- 25 Pearson, J.R., R.C. Pesci and B.H. Iglewski. Submitted for publication.
- Pearson, J.P., K.M. Gray, L. Passador, D.D. Tucker, A. Eberhard, B.H. Igelewski and E.P. Greenberg. 1994. Structure of the autoinducer required for expression of *Pseudomonas aeruginosa* virulence genes. *Proc. Nat. Acad. Sci. USA.* 91:197-201.
- 30 Peterson, G.L. 1997. A simplification of the protein assay method of Lowry et al. which is more generally applicable. *Anal. Biochem.* 83:346-356.

- Schripsema, J., K.E.E. de Rudder, T.G. van Vhet, P.P. Lankhorst, E. de Vroom, J.W. Kijne and A.A.N. van Brussel. 1996. Bacteriocin small of Rhizobium leguminosarum belongs to the class of N-acyl-L-homoserine Lactone molecules, known as autoinducers and as quorum sensing co-transcription factors. J. Bacteriol. 178:366-371.
- Swift, S., M.K. Winson, P.F. Chan, N.J. Bainton, M. Birstall, P.J. Reeves, C.E.C. Rees, S.R. Chhabra, P.J. Hill and G.S.A.B. Stewart. 1993. A novel strategy for the insolation of *lux/homologues* evidence for the widespread distribution of a LuxR:LuxI superfamily in enteric bacteria. Mol. Microbiol. 10:511-520.
- Throup, J., M. Camara, G. Briggs, M.K. Winson, S.R. Chhabra, B.W. Bycroft, P. Williams and G.S.A.B. Stewart. 1995. Characterization of the *yenlyenR* locus from *Yersinia enterocolitica* mediating the synthesis of the N-acylhomoserine lactone signal molecules. Mol. Microbio. 17:345-356.
- Winson, M.K., M. Camara, A. Latifi, M. Foglino, S.R. Chhabra, M. Daykin, M. Bally, V. Chapon, G.P.C. Salmond, B.W. Bycroft, A. Lazdunski, G.S.A.B. Stewart and P. Williams. 1995. Multiple N-acyl-L-homoserine lactone signal molecules regulate production of virulence determinants and secondary metabolites in *Pseudomonas aeruginosa*. Proc. Natl. Acad. Sci. USA. 92:9427-9431.
- Zhang, L., P.J. Murphy and I.T. Max. 1993. Agrobacterium conjugation and gene regulation by N-acyl-J-homoserine lactones. Nature. 362:446-448.
- Allison, D.G. and I.W. Sutherland. 1984. A staining technique for attached bacteria and its correlation to extracellular carbohydrate production. J. Microbiol. Methods. 2:93-99.
- Allison, D.G. and I.W. Sutherland. 1987. The role of polysaccharides in adhesion of freshwater bacteria. J. Gen. Microbiol. 133:1319-1327.
- Anwar, H., J.L. Strap, K. Chen and J.W. Costerton. 1992. Dynamic interactions of biofilms on mucoid *Pseudomonas aeruginosa* with tobramycin and piperacillin. Antimicrob. Agents Chemother. 36:1208-1214.
- Boivin, J. and J.W. Costerton. 1991. Biofilms and biodeterioration. In Biodeterioration and biodegradation 8 (ed.) H.W. Rossmore, p. 53-62. Elsevier Appl. Sci., London.

- Boyd, A., M. Ghosh T.B. May, D. Shinabarger, R. Keogh and A.M. Chakrabarty. 1993. Sequence of the *algL* gene of *Pseudomonas aeruginosa* and purification of its alginate lyase product. *Gene* 131:1-8.
- 5 Boyd, A. and A.M. Chakrabarty. 1994. Role of the alginate lyase in cell attachment of *Pseudomonas aeruginosa*. *Appl. Environ. Microbiol.* 60:2325-2359.
- Boyd, A., R.W. Ye and A.M. Chakrabarty. 1995. Purification and characterization of the
10 alginate lyase of *Pseudomonas aeruginosa*. *J. Bacteriol.* Submitted.
- Christensen, B.E. and W.G. Characklis. 1990. Physical and chemical properties of biofilms. pp. 93-130, *In*: W.G. Characklis and K.C. Marshall (eds.), *Biofilms*. John Wiley & Sons, New York.
- 15 Costerton, J.W., K.J. Cheng and G.G. Gessey. 1987. Bacterial biofilms in nature and disease. *Ann. Rev. Microbiol.* 41:435-464.
- Davies, D.G., A.M. Chakrabarty and G.G. Gessey, 1993. Exopolysaccharide production
20 in biofilms. Substratum activation of alginate gene expression by *Pseudomonas aeruginosa*. *Appl. Environ. Microbiol.* 59:1181-1186.
- Davies, D.G. and G.G. Gessey. 1995. Regulation of the alginate biosynthesis gene *algC* in *Pseudomonas aeruginosa* during biofilm development in continuous culture. *Appl.*
25 *Environ. Microbiol.* 61:860-867.
- Davies, D.G. 1996. Regulation of alginate biosynthesis in *Pseudomonas aeruginosa* biofilms. Doctoral Thesis. Montana State University, Bozeman, MT.
- 30 Dempsey, M.J. 1981. Marine bacterial fouling: a scanning electron microscope study. *Marine Biol.* 61:305-315.
- Fletcher, M. 1980. Adherence of marine micro-organisms to smooth surfaces, pp. 347-371. *In* E.H. Beachey (ed.), *Bacterial Adherence (Receptors and Recognition, series 3,*
35 *vol. 6)*. Chapman & Hall, London.

- Floodgate, G.D., 1972. The mechanism of bacterial attachment to detritus in aquatic systems. *Memorie dell'Istituto italiano di idrobiologica Dott. Carlo di Marchi* 29 (suppl.), 309-323.
- 5 Gacesa, P. 1987. Alginate-modifying-enzymes. A proposed unified mechanism of action for the lyases and epimerases. *FEBS Lett.* 212:199-202.
- Gambello, M.J., S. Kaye and B.H. Iglewski. 1993. LasR of *Pseudomonas aeruginosa* is a transcriptional activator of the alkaline protease gene (*apr*) and an enhancer of
10 exotoxin. A expression. *Infect. Immun.* 61:1180-1184.
- Geesey, G.G., W.T. Richardson, H.G. Yeomans, R.T. Irvin and J.W. Costerton, 1997. Microscopic examination of natural sessile bacterial populations from Alpine streams. *Can. J. Microbiol.* 23:1733-1736.
- 15 Holloway, B.W. 1955. Genetic recombination in *Pseudomonas aeruginosa*. *J. Gen. Microbiol.* 13:572-581.
- Hengge-Atonis, R. 1993. Survival of hunger and stress: the role of *rpoS* in early
20 stationary phase regulation in *Escherichia coli*. *Cell.* 72:165-168.
- Jones, H.C., L.L. Roth and W.M. Sanders. 1969. Electron microscopic study of a slime layer. *J. Bacteriol.* 99:316-325.
- 25 Khoury, A.E., K. Lam, B.D. Ellis and J.W. Costerton. 1992. Prevention and control of bacterial infections associated with medical devices. *ASAIO J.* 38:M174-178.
- Latifi, A., K.M. Winson, M. Foglino, B.S. Bycroft, G.S.A.B. Stewart, A. Lazdunski and P. Williams. 1995. Multiple homologues of LuxR and LuxI control expression of
30 virulence determinants and secondary metabolites through quorum sensing in *Pseudomonas aeruginosa* PAO1. *Mol. Microbiol.* 17:333-344.
- Latifi, A., M. Foglino, K. Tanaka, P. Williams and A. Lazdunski. 1996. A hierarchical quorum-sensing cascade in *Pseudomonas aeruginosa* links the transcriptional activators
35 LasR and RhIR (VsmR) to expression of the stationary-phase sigma factor RpoS. *Mol. Microbiol.* 21:1137-1146.

- Nicholas, W.W., M.J. Evans, M.P.E. Slack and H.L. Walmsley. 1989. The penetration of antibiotics into aggregates of mucoid and nonmucoid *Pseudomonas aeruginosa*. J. Gen. Microbiol. 135:1291-1303.
- 5 Ochsner, U.A. and J. Reiser. 1995. Autoinducer-mediated regulation of rhamnolipid biosurfactant synthesis in *Pseudomonas aeruginosa*. Proc. Natl. Acad. Sci. USA. 92:6424-6428.
- Pearson, J.P., L. Pasador, B.H. Iglewski and E.P. Greenberg. 1995. A second N-
10 acylhomoserine lactone signal produced by *Pseudomonas aeruginosa*. Proc. Natl. Acad. Sci. USA. 92:1490-1494.
- Preiss, J. and G. Ashwell. 1962a. Alginic acid metabolism in bacteria. I. Enzymatic formation of unsaturated oligosaccharides and L-deoxy-L-erythro-5-hexoseulose uronic
15 acid. M. Biol. Chem. 237:309-316.
- Preiss, J. and G. Ashwell. 1962b. Alginic acid metabolism in bacterial. II. The enzymatic reduction of 4-deoxy-1-erythro-5-hexoseulose uronic acid to 2-keto-3-deoxy-D-gluconic acid. J. biol. Chem. 237-321.
20
- Schiller, N.L., S.R. Monday, C.M. Boyd, N.T. Keen and D.E. Ohman. 1993. Characterization of the *Pseudomonas aeruginosa* alginate lyase gene (*algL*): cloning, sequencing and expression in *Escherichia coli*. J. Bacteriol. 175:4780-4789.
- 25 Stinivasan, R., P.S. Stewart, T. Griebel, C.I. Chen and X. Xu. 1995. Biofilm parameters influencing biocide efficacy. Biotech. Bioeng. 46:553-560.
- Stewart, P.S. 1994. Biofilm accumulation model that predicts antibiotic resistance of *Pseudomonas aeruginosa* biofilms. Antimicrob. Agents Chemother. 38:1052-1058.
30
- Sutherland, I.W. 1980. Polysaccharides in the adhesion of marine and freshwater bacteria. pp. 329-338. In R.C. W. Berkeley, J.M. Lynch, J. Melling, P.R. Rutter and B. Vincent (eds.), Microbial Adhesion to Surfaces. Ellis Horwood, London.
- 35 Tachiro, H., T. Numakura, S. Nishikawa and Y. Miyaji. 1991. Penetration of biocides into biofilms. Wat. Sci. Technol. 23:1395-1403.

- 31 -

- Wallace, W.H., J.T. Fleming, D.C. White and G.S. Sayler. 1994. An *algD*-
Bioluminescent reporter plasmid to monitor alginate production in biofilms. *Microb.
Ecol.* 27:225-239.
- 5 Wardell, J.N., C.M. Brown and B. Flannigan. 1983. *In* Microbes and surfaces. Symposia
for the Society for General Microbiology. 34:351-378.
- Winson, M.K., M. Camara, A. Laifi, M. Foglino, S.R. Chhabra, M. Daykin, M. Bally,
V. Chapon, G.P.C. Salmond. B.W. Bycroft, A. Lazdunski, G.S.A.B. Stewart and
10 P. Williams. 1995. Multiple N-acyl-L-homoserine lactone signal molecules regulate
production of virulence determinants and secondary metabolites in *Pseudomonas
aeruginosa*. *Proc. Natl. Acad. Sci. USA.* 92:9427-9431.
- Zobell, C.E. 1943. the effect of solid surfaces upon bacterial activity. *J. Bacteriol.*
15 46:39-56.

The purpose of the above description and examples is to illustrate some
embodiments of the present invention without implying any limitation. It will be
apparent to those of skill in the art that various modifications and variations may be
20 made to the composition and method of the present invention without departing from the
spirit or scope of the invention. All patents and publications cited herein are
incorporated by reference in their entireties.

- 32 -

Claims:

1. A method of regulating biofilm development comprising administering a composition comprising a homoserine lactone compound selected from the group
5 consisting of N-(3-oxododecanoyl) L-homoserine lactone or its analogs, and blocking compounds thereof and butyryl L-homo-serine lactone or its analogs.
2. The method of claim 1 wherein biofilm development is enhanced and stimulated by the addition of N-(3-oxododecanoyl) L-homoserine lactone or its analogs
10 to growing bacterial cultures.
3. The method of claim 1, wherein biofilm development is prevented by the addition of N-(3oxododecanoyl) L-homoserine lactone blocking compounds to growing bacterial cultures.
15
4. The method of claim 1, wherein said biofilm is caused by an integrated bacterial community ($>10^5$) aggregate bacteria.
5. The method of claim 1, wherein detachment and dispersion of bacterial
20 cells in a biofilm is enhanced or stimulated by butyryl L-homoserine lactone.
6. The method of claim 1 wherein detachment and dispersion of bacterial cells in a biofilm can be enhanced or stimulated by the addition of butyryl L-homoserine lactone or by the addition of analogs of butyryl L-homoserine lactone.
25
7. A method for prevention of biofilm matrix polymer development comprising administering a homoserine lactone compound selected from the group consisting of butyryl L-homoserine lactone or its analogs.
8. The method of claim 7, wherein said biofilm matrix polymer is selected
30 from a polypeptide or polysaccharide.
9. A method of biofilm dispersion comprising administering a homoserine lactone compound selected from the group consisting of butyryl L-homoserine lactone
35 or its analogs.

- 33 -

10. A method of prevention of biofilm dispersion comprising administering an effective amount of a composition including N-(3-oxododecanoyl) L-homoserine lactone or its analogs.
- 5 11. The method of claim 1, wherein said biofilm is a product of bacteria.
12. The method of claim 1, wherein said biofilms are mixed biofilms.
13. The method of claim 12 wherein said mixed biofilm is the product of two
10 or more of the group consisting of bacteria, fungi and protozoa.
14. The method of claim 1, wherein said composition further comprises a surfactant.
- 15 15. The method of claim 14 wherein said surfactant is selected from the group consisting of Sodium Dodecyl Sulfate; Quaternary Ammonium Compounds; alkyl pyridinium iodides; Tween 80, Tween 85, Triton X-100; Brij 56; Biological surfactants; Rhamnolipid, Surfactin, and Visconsin, and sulfonates.
- 20 16. The method of claim 1, wherein said method further comprises dispersion of the biofilm with 0.2% or higher SDS.
17. The method of claim 1, wherein said biofilms comprise a member of the group consisting of polyuronic acids, polysaccharides, polypeptides, alginates and
25 mixtures thereof.
18. The method of claim 7, wherein said composition further comprises 3-oxo-N-(tetrahydro-2-oxo-3-furanyl) hexanamide.
- 30 19. A composition for regulating biofilm development comprising a homoserine lactone compound selected from the group consisting of N-(3-oxododecanoyl) L-homoserine lactone or its analogs, blocking compounds and butyryl L-homo-serine lactone or its analogs, and a carrier.
- 35 20. The composition of claim 19, wherein said biofilm is a product of bacteria.

- 34 -

21. The composition of claim 20, wherein said biofilm is a mixed biofilm.

22. The composition of claim 21 wherein said mixed biofilm is the product of two or more of the group consisting of bacteria, fungi and protozoa.

5

23. The composition of claim 19, wherein said composition further comprises a surfactant.

24. The composition of claim 23 wherein said surfactant is selected from the group consisting of Sodium Dodecyl Sulfate; Quaternary Ammonium Compounds; alkyl
10 pyridinium iodides; Tween 80, Tween 85, Triton X100; Brij 56; Biological surfactants; Rhamnolipid, Surfactin, and Visconsin, and sulfonates.

25. The composition of claim 19 wherein said composition further comprises
15 dispersion of the biofilm with 0.2% or higher SDS.

26. The composition of claim 19 wherein said biofilms comprise a member of the group consisting of polyuronic acids, polysaccharides, polypeptides, alginates and mixtures thereof.

20

27. A method of cleaning a surface which comprises administering a composition comprising a homoserine lactone compound selected from the group consisting of N(3-oxododecanoyl) L-homoserine lactone or its analogs and blocking compounds, and butyryl L-homo-serine lactone or its analogs.

25

28. The method of claim 27 wherein biofilm development is enhanced and stimulated by the addition of N-(3-oxododecanoyl) L-homoserine lactone or its analogs to growing bacterial cultures.

29. The method of claim 27, wherein biofilm development is prevented by
30 the addition of N-(3-oxododecanoyl) L-homoserine lactone blocking compounds to growing bacterial cultures.

30. The method of claim 27, wherein said biofilm is caused by an integrated
35 bacterial community ($>10^5$) aggregate bacteria.

- 35 -

31. The method of claim 27, wherein detachment and dispersion of bacterial cells in a biofilm is enhanced or stimulated by butyryl L-homoserine lactone.

32. The method of claim 27 wherein detachment and dispersion of bacterial
5 cells in a biofilm can be enhanced or stimulated by the addition of butyryl L-homoserine lactone or by the addition of analogs of butyryl L-homoserine lactone.

33. A method according to claim 27 wherein the surface to be cleaned is a
10 hard surface, woven surface, or non-woven surface.

34. A method according to claim 27 wherein the surface to be cleaned is a
toilet bowl, bath tub, drain, chair, countertop, food surfaces, airduct, air conditioner,
carpet, paper, or cloth.

35. A topical dressing for burn patients comprising a homoserine lactone
15 compound selected from the group consisting of butyryl L-homoserine lactone or its analogs.

36. A dentifrice comprising a homoserine lactone compound selected from
20 the group consisting of blockers for the group of molecules consisting of OdDHL, from the group consisting of butyryl L-homoserine lactone or its analogs.

37. A mouthwash comprising a homoserine lactone compound selected from
the group consisting of butyryl L-homoserine lactone or its analogs or an N-(3-oxo-
25 decanoyl) L-homoserine lactone blocking compound.

38. A method for treatment and prevention of dental caries comprising
administering a composition comprising an effective amount of homoserine lactone
compound selected from the group consisting of butyryl L-homoserine lactone or its
30 analogs, or an OdDHL blocking compound.

39. A method for the treatment of acne comprising administering a
composition comprising an effective amount of homoserine lactone compound selected
from the group consisting of butyryl L-homoserine lactone or its analogs, or an OdDHL
35 blocking compound.

- 36 -

40. A method for cleaning and disinfecting contact lenses comprising administering a composition comprising an effective amount of homoserine lactone compound selected from the group consisting of butyryl L-homoserine lactone or its analogs.

5

41. A method of treating medical indwelling devices comprising administering a composition comprising an effective amount of homoserine lactone compound selected from the group consisting of butyryl L-homoserine lactone or its analogs or an N-(3-oxodo-decanoyl) L-homoserine lactone blocking compound.

10

42. The method of claim 44, wherein said device is selected from the group consisting of catheters, orthopedic devices and implants.

43. The composition of claim 35 wherein said composition includes a topical cream, transdermal, ointment or oil.

15

44. A microbial disinfectant composition comprising an effective amount of homoserine lactone compound selected from the group consisting of butyryl - homoserine lactone or its analogs.

20

45. The disinfectant of claim 44, wherein said microbe is selected from the group consisting of fungi, protozoa and gram negative bacteria.

46. The disinfectant of claim 45 wherein said gram negative bacteria are selected from the group consisting of *Pseudomonadaceae*, *Azotobacteraceae*, *Rhizobiaceae*, *Methylococcaceae*, *Halobacteriaceae*, *Acetobacteraceae*, *Legionellaceae*, *Neisseriaceae* and other Genera.

25

47. The disinfectant of claim 44, further comprising a biocide or antibiotic.

30

48. The disinfectant of claim 44, wherein said disinfectant is applied to drains, shower curtains, grout, toilets, flooring.

49. The disinfectant of claim 44, wherein said disinfectant is in a formulation selected from the group consisting of a spray, powder and liquid formulation.

35

- 37 -

50. The disinfectant of claim 49, wherein said bacteria is of the class *Pseudomonas*.

51. The disinfectant of claim 50, wherein said bacteria is of the species
5 *Pseudomonas aeruginosa*.

52. A method of blocking the binding interaction of homoserine lactones with their corresponding DNA binding protein products of LasI and Las R comprising administering a homoserine lactone blocking compound, wherein said blocking
10 interaction prevents biofilm formation.

53. A method of preventing fouling comprising administering a composition comprising a homoserine lactone compound selected from the group consisting of N-(3-oxododecanoyl) L-homoserine lactone blocking compounds and butyryl L-homoserine
15 lactone or its analogs.

54. The method of claim 53 wherein said fouling is in a system selected from the group consisting of injection wells for oil recovery, cooling towers, water purification systems, porous media (soil, sand), marine environments and hospital or
20 automotive airconditioning systems.

55. A method for the dispersal of environmental concentrations of bacteria comprising administering a homoserine lactone compound selected from the group consisting of N-(3-oxododecanoyl) L-homoserine lactone blocking compounds and
25 butyryl L-homoserine lactone or its analogs to the environmental concentration of bacteria.

56. A disinfectant coating comprising a homoserine lactone compound selected from the group consisting of N(3-oxododecanoyl) L-homoserine lactone blocking compounds and butyryl L-homoserine lactone or its analogs copolymerized to a
30 polymer.

57. The disinfectant coating of claim 56, wherein said disinfectant is applied to drains, shower curtains, grout, toilets, flooring.

- 38 -

58. A method of enhancing biofilm formation in fermentation comprising adding N-(3-oxodo-decanoyl) L-homoserine lactone or its analogs to a fermentation system.


5 59. A specific therapy to reduce gram negative bacterial without depleting beneficial gram positive bacteria comprising N-(3-oxododecanoyl) L-homoserine lactone blocking compounds and butyryl L-homoserine lactone or its analogs.

60. The method of claim 59 wherein said therapy is for the treatment of a
10 condition selected from the group consisting of urethral E. coli infections and vaginal candida infections.

61. A method of treating a condition selected from the group consisting of middle ear infections (children), osteomyelitis and prostatitis comprising administering a
15 homoserine lactone compound selected from the group consisting of N-(3-oxododecanoyl) L-homoserine lactone blocking compounds and butyryl L-homoserine lactone or its analogs, wherein said homoserine lactone compound disperses bacteria making them more susceptible to antibiotics.


1/15

Fig 1




2/15

Fig 2




3/15

Fig3.



4/15

Fig 4




5/15



Figure 5


6/15

Figure 6




7/15

Figure 7




8/15

Figure 8



9/15

Figure 9




10/15

Figure 10



11/15


Figure 11



12/15

Figure 12

Addition of 0.2% Sodium Dodecyl Sulfate
to Cell Clusters of
P. aeruginosa PAO1 JP2 and *P. aeruginosa* PAO1 parent




Panel B

Panel A

13/15

Figure 13


Addition of 0.2% SDS to cell cluster of
Pseudomonas aeruginosa JP2
grown in spent medium from parental strain



14/15

Figure 14

↙ void space



Exterior edge
of aggregate ↗

Figure 15

BHL

Effect of Addition of to *P. aeruginosa* PDO100 biofilm



Biofilm before addition of factor 2



Biofilm 18 hours after addition of
factor 2 (higher magnification of the biofilm)